



ELSEVIER

Available online at www.sciencedirect.com

# MAPK cascade signalling networks in plant defence

Andrea Pitzschke<sup>1,3</sup>, Adam Schikora<sup>2,3</sup> and Heribert Hirt<sup>1,2</sup>

The sensing of stress signals and their transduction into appropriate responses is crucial for the adaptation and survival of plants. Kinase cascades of the mitogen-activated protein kinase (MAPK) class play a remarkably important role in plant signalling of a variety of abiotic and biotic stresses. MAPK cascade-mediated signalling is an essential step in the establishment of resistance to pathogens. Here, we describe the most recent insights into MAPK-mediated pathogen defence response regulation with a particular focus on the cascades involving MPK3, MPK4 and MPK6. We also discuss the strategies developed by plant pathogens to circumvent, inactivate or even 'hijack' MAPK-mediated defence responses.

## Addresses

<sup>1</sup> Max F. Perutz Laboratories, University of Vienna, Dr. Bohrgasse 9, 1030 Vienna, Austria

<sup>2</sup> URGV Plant Genomics, INRA-CNRS-University of Evry, 2 rue Gaston Cremieux, 91057 Evry, France

Corresponding author: Hirt, Heribert ([hirt@evry.inra.fr](mailto:hirt@evry.inra.fr))

<sup>3</sup> Both authors contributed equally to this work.

Current Opinion in Plant Biology 2009, 12:1–6

This review comes from a themed issue on  
Biotic Interactions  
Edited by Xinnian Dong and Regine Kahmann

1369-5266/\$ – see front matter  
© 2009 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.pbi.2009.06.008

## Introduction

MAPK cascades are highly conserved modules in all eukaryotes. In plants, MAPK pathways are involved in the regulation of development, growth, programmed cell death and in responses to a diversity of environmental stimuli including cold, heat, reactive oxygen species, UV, drought and pathogen attack [1]. Via a phosphorelay mechanism these cascades, minimally composed of a MAPKKK (MAPK kinase kinase), a MAPKK (MAPK kinase) and a MAPK, link upstream receptors to downstream targets. The *Arabidopsis* genome contains about 110 genes coding for putative MAPK pathway components: 20 MAPKs, 10 MAPKKs and more than 80 MAPKKKs [2]. Scaffolding proteins or specific expression of distinct components, confer the specificity to MAPK components engaged in a module. So far, only a few MAPK cascade components have been studied in detail. The *Arabidopsis* MAPKKKs YODA, ANP2/ANP3 and

MP3K6/MP3K7 function in development [3<sup>••</sup>,4–6], MEKK1 and ANP1 act in the environmental stress response [7<sup>•</sup>–9<sup>•</sup>], whereas CTR1 plays a pivotal role in ethylene signalling [10]. Among the 10 MAPKKs, MKK1/MKK2, MKK4/MKK5 as well as MKK3, MKK7 and MKK9 have been analyzed [11,12,13<sup>••</sup>,14<sup>••</sup>,15,16<sup>•</sup>,17,18<sup>••</sup>,19,20<sup>••</sup>]. Finally, 8 of the 20 MAPKs have been studied to various degrees (for review, see [1]).

The best-characterized MAPKs are MPK3, MPK4 and MPK6, all of which are activated by a diversity of stimuli including abiotic stresses, pathogens and oxidative stress. While MPK4 negatively regulates biotic stress signalling, MPK3 and MPK6 act as positive mediators of defence responses. The key role of these three MAPKs for normal plant growth and development is evidenced by the severely dwarfed phenotype of *mpk4* and the embryo lethal phenotype of *mpk3/mpk6* double mutants [14<sup>••</sup>,16<sup>•</sup>,21].

In this review, we focus on the activation of MAPK cascades involving MPK3, MPK4 and MPK6, after the perception of pathogen-associated molecular patterns (PAMPs), and on strategies developed by specific plant pathogens to deal with MAPK-mediated defence responses.

## PAMP-triggered activation of MAPK cascades

In the ongoing battle between plants and pathogens, plants have adapted the capacity to recognize pathogens through PAMPs via cell surface-located pathogen-recognition receptors. The activation of these receptors induces convergent intracellular signalling pathways in plant cells, which ultimately result in the establishment of PAMP-triggered immunity [22]. PAMPs are mostly small molecules derived from different pathogen structures common to a class of pathogens. Consequently, the responses to PAMPs are not specific, but rather reflect a response to a given group of pathogens. Despite the large variety of known PAMPs, PAMP receptors in plants have so far only been identified for flagellin, the translation elongation factor EF-Tu and chitin [23,24<sup>••</sup>,25,26,27<sup>••</sup>].

PAMP-triggered immunity requires a signal transduction from receptors to downstream components via the MAPK cascade, and many of the known PAMPs were shown to activate MAP kinases. The flagellin derived peptide flg22 triggers a rapid and strong activation of MPK3, MPK4 and MPK6 [28]. MPK4 and MPK6 are also activated by harpin proteins, which are encoded by *hrp* (hypersensitive response and pathogenicity) genes in

## 2 Biotic Interactions

many plant pathogenic bacteria. This activation is followed by the induction of pathogenesis-related (*PR*) genes [29], encoding for proteins with antimicrobial activities. Similarly, various NLPs (necrosis and ethylene-inducing peptide1-like proteins) trigger MAPK activation and induce defence responses [30].

### MPK3/MPK6 are necessary to induce defence responses

MPK3 and MPK6 are closely related proteins that show a high level of functional redundancy. Both MAPKs are key regulators of a diverse set of processes including abscission, stomatal development, signal various abiotic stresses and defence response to bacterial and fungal pathogens (for review see [1]).

On the basis of experiments using transient expression in protoplasts, the MAPK module MEKK1-MKK4/MKK5-MPK3/MPK6 was proposed to be responsible for flg22 signal transmission [20<sup>•</sup>]. The involvement of MEKK1 in flg22-induced MKK4/MKK5-MPK3/MPK6 signalling is unlikely, since *mekk1* mutant plants are compromised in flg22-triggered activation of MPK4, but show normal activation of MPK3 and MPK6 [9<sup>•</sup>]. However, besides this uncertainty, modules involving the MAPKKs MKK1/MKK4/MKK5/MKK9 and the MAPKs MPK3/MPK6 are clearly implicated in different defence strategies.

### MPK3/MPK6 are compulsory in camalexin biosynthesis

Intensive research has been attributed to plant defence responses to *Botrytis cinerea*. This fungal pathogen triggers the synthesis of camalexin, a major phytoalexin in *Arabidopsis*. Camalexin is required for resistance to *B. cinerea*, as shown by the susceptible phenotype of the *phytoalexin deficient 3 (pad3)* mutant [32]. The key role of MPK3/MPK6 in camalexin-based fungal resistance is demonstrated by the fact that *mpk3* and *mpk6* mutants are compromised in camalexin production and consequently more susceptible to *B. cinerea* [31]. Recent data implicate a MAPK cascade composed of MKK4/MKK5 and MPK3/MPK6 in response to fungal pathogens, based on the observation that activation of MPK3/MPK6 in conditional gain-of-function (GOF) plants for MKK4/MKK5 or MEKK1/MKKKa is sufficient to induce accumulation of camalexin, even in the absence of pathogen attack [31]. However, another MAPKK, MKK9, whose upstream MAPKKK is unknown, is apparently also involved in camalexin biosynthesis via MPK3/MPK6: Conditional MKK9-GOF plants have elevated MPK3/MPK6 activity and produce even more camalexin after transgene induction than MKK4-GOF or MKK5-GOF plants [17]. In addition, the MKK9-MPK3/6 is involved in the biosynthesis of ethylene [17] a plant hormone involved in defence responses. Whether both MKK4/MKK5 and MKK9 act as upstream regulators of MPK3/MPK6 in camalexin biosynthesis will have to be verified.

### MPK3 is required for stomatal immune responses

Stomata are indispensable for gas exchange and transpiration. However, in terms of pathogen attack stomata are weak points, since they serve as 'entrance gate' for various microbial invaders. A tight regulation of stomatal opening and closure, therefore, is pivotal not only to withstand unfavourable environmental conditions, such as drought and heat but also to restrict pathogen invasion. Recent studies implicate MAPK cascades in stomatal regulation in both abiotic and biotic stress responses.

MPK3/MPK6 are key players in stomatal development and stomatal dynamics [16<sup>•</sup>]. Upon drought, stomatal closure is mediated through the phytohormone ABA and involves MKK1, MPK3 and MPK6 [33,34]. Pathogen-induced stomatal closure restricts the invasion of many bacteria and is a part of the plant innate immune response. The pathogen *Xanthomonas campestris* pv. *Campestris* (*Xcc*) excretes a cell-cell signal-regulated virulence factor, which reverts stomatal closure induced by bacteria and ABA [33] and which can complement the infectivity of *Pseudomonas syringae* pv. *tomato* (*Pst*) mutants deficient in the production of coronatine, a toxin required to overcome stomatal defence. Recent findings suggest a unique function of MPK3 in the stomatal innate immunity response. Guard cell specific Arabidopsis *MPK3* antisense plants are more sensitive to *Pst* coronatine-deficient mutants. Their stomata close normally upon ABA [33], but are unresponsive to bacteria. Moreover, in these plants, *Xcc* extracts do not revert bacteria-induced or ABA-induced stomatal closure. Whether pathogen-induced and ABA-induced stomatal closures are signalled via a common MAPK cascade remains to be investigated.

### Negative regulation of defence responses by the MPK4 pathway

A number of studies identified the MEKK1-MKK1/2-MPK4 cascade in pathogen signalling. Using genetic approaches, independent reports show MKK1 and MKK2 as functionally redundant activators of MPK4 [13<sup>••</sup>,14<sup>••</sup>,35<sup>••</sup>], thus confirming previous experimental evidence that MKK1 and MKK2 interact with MPK4 [36,37].

*Mekk1, mkk1/mkk2* double and *mpk4* mutants are severely dwarfed and accumulate high amounts of reactive oxygen species [7<sup>•</sup>-9<sup>•</sup>,13<sup>••</sup>,14<sup>••</sup>,38]. These abnormalities are most probably due to their drastically enhanced SA levels and can be partially reverted by expression of a bacterial SA hydrolase [38]. *Mekk1, mkk1/mkk2* double and *mpk4* mutants also display spontaneous cell death, upregulation of pathogenesis-related genes and enhanced resistance to pathogens. Therefore, a negative regulatory role of the MEKK1-MKK1/2-MPK4 module in SA and H<sub>2</sub>O<sub>2</sub> production has been proposed.

The *mekk1*, *mkk1/2* and *mpk4* mutants have strongly overlapping expression profiles, with many defence-related genes being similarly deregulated [14<sup>••</sup>]. Interestingly, comparative transcriptome studies have implicated that a subset of the MEKK1-dependent MPK4 responses are regulated independently of MKK1 and MKK2 [35<sup>••</sup>]. In addition, based on the significant proportion of genes exclusively deregulated in *mekk1* mutants, MEKK1 seems to function in other as yet unknown pathways involving neither MKK1/MKK2 nor MPK4. The WRKY53 transcription factor may be partially responsible for the *mekk1*-specific gene set, as MEKK1 directly interacts with WRKY53 and alters the activity of this transcription factor [39<sup>•</sup>—a unique short cut of MAPK signalling. The interaction of MKK1-MPK4 and MKK2-MPK4 was lately confirmed *in vivo*, and the complexes were located at the plasma membrane and in the nucleus [13<sup>••</sup>]. Three proteins have been identified as substrates of MPK4: WRKY33, WRKY25 and MKS1 [40]. WRKY33 was previously shown to interact with MKS1 *in yeast* and *in vivo* [38]. Although no MPK4-WRKY33 interaction was originally observed, recent results propose an MKS1-dependent MPK4-WRKY33 interaction [41<sup>••</sup>]. Immunoprecipitation from nuclear extracts revealed the existence of a ternary MKS1-MPK4-WRKY33 complex. The recruitment of WRKY33 to this complex depends on the phosphorylation state of MPK4. After activation, MPK4 multiphosphorylates MKS1, resulting in the release of WRKY33 from the ternary complex. According to this model, free WRKY33 then induces transcription of its target genes [41<sup>••</sup>], providing a molecular mechanism for the negative regulation of defence signalling by MPK4. In the absence of functional MPK4, WRKY33 (and probably other MPK4 targets) would not be sequestered in the complex any more, leading to activation of gene expression even in the absence of pathogens.

### How pathogens manipulate MAPK signalling

Pathogens have evolved strategies to overcome defence responses. This can be achieved through inactivation of PAMP-induced signalling pathways by targeting the MAPK cascade components.

The pathogen *Pseudomonas syringae* injects a number of effectors into plant cells. Among these, AvrPto and AvrPtoB interact with the FLS2 receptor and its co-receptor BAK1. AvrPtoB catalyses the polyubiquitination and subsequent proteasome-dependent degradation of FLS2 [42<sup>••</sup>,43]. This process is enhanced when FLS2 binds to flg22. AvrPto can interact with BAK1 and thereby prevent its binding to FLS2 [42<sup>••</sup>]. In this way, both AvrPto and AvrPtoB interrupt signalling to the downstream MAPK module.

*Pseudomonas syringae* has yet another factor that can directly interact with the MAPK cascade components: HopAI1 is a phosphothreonine lyase that dephosphory-

lates the threonine residue at which MAPKs are activated by their upstream MAPKKs [44<sup>•</sup>]. HopAI1 (when expressed *in planta*) directly interacts with MPK3 and MPK6, thus attenuating flg22-induced MAPK activation and downstream defence responses. Strikingly, HopAI1 is also present in animal/human pathogens such as *Shigella* spp. (OspF) [45,46] and *Salmonella* spp. (SpvC) [47], where it interacts with the MAPKs ERK1/2 and p38.

HopPtoD2 is another bacterial effector from *P. syringae* *pv. tomato* that displays phosphatase activity [48]. Expression of HopPtoD2 in tobacco cells suppresses the cell death provoked by expression of the constitutively active MAPKK variant NtMEK2<sup>DD</sup>. However, HopPtoD2 does not inhibit flg22-induced MAPK activation in *Arabidopsis*.

### *Agrobacterium* hijacks the MPK3/MPK6 pathway


An entirely different mode of host MAPK signalling manipulation by pathogens has recently been unravelled. The soil-borne pathogen *Agrobacterium tumefaciens* carries flagellin variants that are non-detectable by the *Arabidopsis* FLS2 [24<sup>••</sup>,49], yet it triggers a typical innate immune response through the EF-Tu protein. The EF-Tu-derived peptide elf18 is sufficient to trigger the defence response. Interestingly, the receptors for flg22 and elf18, FLS2 and EFR, respectively, belong to the same subfamily of LRR-RLKs, LRRXII. Moreover, elf18 and flg22 induce an overlapping set of responses, including extracellular alkalinisation, a rapid activation of MAPKs and the induction of similar response genes [27<sup>••</sup>]. It is tempting to speculate that also other members of the LRRXII subfamily function as PAMP receptors and it will be interesting to see whether their signalling converge in the same pathways.

*Agrobacterium tumefaciens* is the causal agent of crown galls. It infects plants by integrating a segment of its DNA (transfer DNA) into the host chromosomal DNA. *efr1* mutants fail to recognize EF-Tu and, presumably owing to reduced defence responses, are more easily transformed by *Agrobacterium* [27<sup>••</sup>]. However, since MAPK activities in *efr1* have not been assessed, and since *Agrobacterium* contains other PAMPs in addition to EF-Tu, the importance/contribution of MAPKs in EF-Tu-triggered defence remains unclear.

Initiation of defence signalling cascades in their host can be turned into a benefit for plant pathogens: The activation of MPK3 in response to flg22 or *Agrobacterium* results in the phosphorylation and subsequent nuclear translocation of the host protein VIP1 (virE2 interacting protein 1). *Agrobacterium* has hijacked VIP1 for delivering their T-DNA into the plant nucleus, where it integrates into the host genome [50,51<sup>••</sup>]. Because VIP1 does not only serve as nuclear shuttle for the pathogenic T-DNA complex but can also induce the expression of defence

## 4 Biotic Interactions

Figure 1



PAMP-induced MAPK cascades in the plant defence to bacterial and fungal pathogens. PAMP-triggered activation of MAPK cascades initiates the synthesis of pathogen-induced synthesis of PR proteins (e.g. glucanases, chitinases), cell wall depositions, stomatal closure and phytoalexin (e.g. camalexin) synthesis. Together, these modifications result in pathogen resistance. Microbial factors interfering with the signal transduction are shown in blue. Activated transcription factors (TF) are shown as semicircles. Unknown receptors and MAPK cascade components are indicated by '?'.

genes [51<sup>••</sup>], nuclear VIP1 would be counteracting *Agrobacterium* invasion. *Agrobacterium* overcomes this problem through targeting nuclear VIP1 for proteasome degradation by the *Agrobacterium* virulence factor VirF, which encodes an F-box protein [52].

## Conclusions

By amplifying and transducing pathogen-derived signals perceived at membrane receptors and transducing these signals into altered gene expression, plant MAPK modules play a key role in the induction of defence mechanisms (Figure 1).

MAPKs are also prominent targets for inactivation by effector proteins. Interestingly, the mechanisms of effector-mediated interruption of MAPK signalling employed by plant and animal pathogens are similar. So far only a few components of MAPK cascades have been thoroughly studied in plants. Three reports described

the systematic attempts to interconnect the MAPKK and MAPKs with their targets [53,54<sup>•</sup>,55]. The identification and *in planta* verification of additional MAPKKK-MAPKK-MAPK modules, discovery of downstream targets and the regulated processes will be a future challenge to disentangle the sophisticated network of plant MAPK signalling pathways.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Colcombet J, Hirt H: **Arabidopsis MAPKs: a complex signalling network involved in multiple biological processes**. *Biochem J* 2008, **413**:217-226.
  2. MAPK G, Ichimura K, Shinozaki K, Tena G, Sheen J, Henry Y, Champion A, Kreis M, Zhang S, Hirt H *et al.*: **Mitogen-activated protein kinase cascades in plants: a new nomenclature**. *Trends Plant Sci* 2002, **7**:301-308.

3. Bergmann DC, Lukowitz W, Somerville CR: **Stomatal development and pattern controlled by a MAPKK kinase.** *Science* 2004, **304**:1494-1497.
- The authors describe the MAPKKK YODA as a negative regulator of stomata development. *yoda* null mutants have stomata excess, whereas YODA overexpressing plants lack stomata. Transcriptome analysis of seedlings with altered YODA activity was used to identify potential stomatal regulatory genes. A putative transcription factor was shown to regulate the developmental behaviour of stomatal precursors.
4. Chaiwongsar S, Otegui MS, Jester PJ, Monson SS, Krysan PJ: **The protein kinase genes MAP3K epsilon 1 and MAP3K epsilon 2 are required for pollen viability in Arabidopsis thaliana.** *Plant J* 2006, **48**:193-205.
5. Krysan PJ, Jester PJ, Gottwald JR, Sussman MR: **An Arabidopsis mitogen-activated protein kinase kinase kinase gene family encodes essential positive regulators of cytokinesis.** *Plant Cell* 2002, **14**:1109-1120.
6. Lukowitz W, Roeder A, Parmenter D, Somerville C: **A MAPKK kinase gene regulates extra-embryonic cell fate in Arabidopsis.** *Cell* 2004, **116**:109-119.
7. Ichimura K, Casais C, Peck SC, Shinozaki K, Shirasu K: **MEKK1 is required for MPK4 activation and regulates tissue-specific and temperature-dependent cell death in Arabidopsis.** *J Biol Chem* 2006, **281**:36969-36976.
- Dwarfism and cell death in *mekk1* and *mpk4* mutants is salicylic-acid dependent and can be suppressed by growing the mutants at elevated temperatures. The authors suggest that MEKK1 negatively regulates cell death pathways activated by certain R protein(s) that are unstable at high temperature.
8. Nakagami H, Soukupova H, Schikora A, Zarsky V, Hirt H: **A mitogen-activated protein kinase kinase mediates reactive oxygen species homeostasis in Arabidopsis.** *J Biol Chem* 2006, **281**:38697-38704.
- This paper highlights the commonalities (phenotype, cell death, expression profiles) between *mekk1* and *mpk4* mutants and positions MEKK1 and MPK4 in a common signalling pathway.
9. Suarez-Rodriguez MC, Adams-Phillips L, Liu Y, Wang H, Su SH, Jester PJ, Zhang S, Bent AF, Krysan PJ: **MEKK1 is required for flg22-induced MPK4 activation in Arabidopsis plants.** *Plant Physiol* 2007, **143**:661-669.
- The *mekk1* phenotype was partially rescued by overexpression of a kinase-impaired-version of MEKK1 (K361M), and K361M was sufficient to signal flg22-triggered activation of MPK4; suggesting that MEKK1 may play a structural role in signaling, independent of its protein kinase activity.
10. Kendrick MD, Chang C: **Ethylene signaling: new levels of complexity and regulation.** *Curr Opin Plant Biol* 2008, **11**:479-485.
11. Dai Y, Wang H, Li B, Huang J, Liu X, Zhou Y, Mou Z, Li J: **Increased expression of MAP KINASE KINASE7 causes deficiency in polar auxin transport and leads to plant architectural abnormality in Arabidopsis.** *Plant Cell* 2006, **18**:308-320.
12. Doczi R, Brader G, Pettko-Szandtner A, Rajh I, Djamei A, Pitzschke A, Teige M, Hirt H: **The Arabidopsis mitogen-activated protein kinase kinase MKK3 is upstream of group C mitogen-activated protein kinases and participates in pathogen signaling.** *Plant Cell* 2007, **19**:3266-3279.
13. Gao M, Liu J, Bi D, Zhang Z, Cheng F, Chen S, Zhang Y: **MEKK1, MKK1/MKK2 and MPK4 function together in a mitogen-activated protein kinase cascade to regulate innate immunity in plants.** *Cell Res* 2008, **18**:1190-1198.
- New alleles of *mpk4* and *mekk1* were identified. Both MPK4 and MEKK1 were shown to interact with MKK1 and MKK2. *mekk1/mpk4* double mutants were generated. The *mpk4*, *mekk1* and the *mekk1/mpk4* double mutants have similar phenotypes. The impaired activation of MPK4 by flg22 in the *mekk1/2* double mutant suggests that MEKK1, MKK1/MKK2 and MPK4 are in common pathway.
14. Qiu J-L, Zhou L, Yun B-W, Nielsen HB, Fill BK, Petersen K, MacKinlay J, Loake GJ, Mundy J, Morris PC: **Arabidopsis mitogen-activated protein kinases MKK1 and MKK2 have overlapping functions in defense signaling mediated by MEKK1, MPK4, and MKS1.** *Plant Physiol* 2008, **148**:212-222.
- Similar to [13\*]; the authors characterize a *mekk1/mpk4* double mutant. In addition, transcriptome analyses delineate overlapping as well as specific effects of the kinases on global gene expression patterns demonstrating both redundant and unique functions for MKK1 and MKK2.
15. Takahashi F, Yoshida R, Ichimura K, Mizoguchi T, Seo S, Yonezawa M, Maruyama K, Yamaguchi-Shinozaki K, Shinozaki K: **The mitogen-activated protein kinase cascade MKK3-MPK6 is an important part of the jasmonate signal transduction pathway in Arabidopsis.** *Plant Cell* 2007, **19**:805-818.
16. Wang H, Ngwenyama N, Liu Y, Walker JC, Zhang S: **Stomatal development and patterning are regulated by environmentally responsive mitogen-activated protein kinases in Arabidopsis.** *Plant Cell* 2007, **19**:63-73.
- Using various RNA interference and mutant lines the authors provide evidence for a complete MAPK cascade in the regulation of stomatal development and distribution: YODA-MKK4/MKK5-MPK3/MPK6.
17. Xu J, Li Y, Wang Y, Liu H, Lei L, Yang H, Liu G, Ren D: **Activation of MAPK kinase 9 induces ethylene and camalexin biosynthesis and enhances sensitivity to salt stress in Arabidopsis.** *J Biol Chem* 2008, **283**:26996-27006.
18. Yoo SD, Cho YH, Tena G, Xiong Y, Sheen J: **Dual control of nuclear EIN3 by bifurcate MAPK cascades in C2H4 signalling.** *Nature* 2008, **451**:789-795.
- The dual phosphorylation on EIN3 has different outcomes: pT174 promotes stabilization of EIN3 and thereby ethylene responses, while pT592 leads to EIN3 degradation. A new MKK9-MPK3/9 cascade is proposed to be responsible for the ethylene-dependent T174 phosphorylation of EIN3.
19. Zhang X, Dai Y, Xiong Y, DeFraia C, Li J, Dong X, Mou Z: **Overexpression of Arabidopsis MAP kinase kinase 7 leads to activation of plant basal and systemic acquired resistance.** *Plant J* 2007, **52**:1066-1079.
20. Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, Gomez-Gomez L, Boller T, Ausubel FM, Sheen J: **MAP kinase signalling cascade in Arabidopsis innate immunity.** *Nature* 2002, **415**:977-983.
21. Petersen M, Brodersen P, Naested H, Andreasson E, Lindhart U, Johansen B, Nielsen HB, Lacy M, Austin MJ, Parker JE *et al.*: **Arabidopsis map kinase 4 negatively regulates systemic acquired resistance.** *Cell* 2000, **103**:1111-1120.
22. Jones JD, Dangl JL: **The plant immune system.** *Nature* 2006, **444**:323-329.
23. Gaulin E, Drame N, Lafitte C, Torto-Alalibo T, Martinez Y, Ameline-Torregrosa C, Khatib M, Mazarguil H, Villalba-Mateos F, Kamoun S *et al.*: **Cellulose binding domains of a Phytophthora cell wall protein are novel pathogen-associated molecular patterns.** *Plant Cell* 2006, **18**:1766-1777.
24. Gomez-Gomez L, Felix G, Boller T: **A single locus determines sensitivity to bacterial flagellin in Arabidopsis thaliana.** *Plant J* 1999, **18**:277-284.
25. Kaku H, Nishizawa Y, Ishii-Minami N, Akimoto-Tomiya C, Dohmae N, Takio K, Minami E, Shibuya N: **Plant cells recognize chitin fragments for defense signaling through a plasma membrane receptor.** *Proc Natl Acad Sci U S A* 2006, **103**:11086-11091.
26. Livaja M, Zeidler D, von Rad U, Durner J: **Transcriptional responses of Arabidopsis thaliana to the bacteria-derived PAMPs harpin and lipopolysaccharide.** *Immunobiology* 2008, **213**:161-171.
27. Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JD, Boller T, Felix G: **Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts Agrobacterium-mediated transformation.** *Cell* 2006, **125**:749-760.
- Identification of the EF-Tu receptor. This report shows that the reactions to elf18 and flg22 significantly overlap. The authors suggest that both PAMPs induce the same signalling pathway. An *efr* mutant is more susceptible to Agrobacterium transformation, indicating that PAMP-induced defence mechanisms may interfere with transformation process.
28. Droillard MJ, Boudsocq M, Barbier-Brygoo H, Lauriere C: **Involvement of MPK4 in osmotic stress response pathways in cell suspensions and plantlets of Arabidopsis thaliana: activation by hypoosmolarity and negative role in hyperosmolarity tolerance.** *FEBS Lett* 2004, **574**:42-48.

## 6 Biotic Interactions

29. Desikan R, Hancock JT, Ichimura K, Shinozaki K, Neill SJ: **Harpin induces activation of the arabidopsis mitogen-activated protein kinases AtMPK4 and AtMPK6.** *Plant Physiol* 2001, **126**:1579-1587.
30. Qutob D, Kemmerling B, Brunner F, Kufner I, Engelhardt S, Gust AA, Luberacki B, Seitz HU, Stahl D, Rauhut T *et al.*: **Phytotoxicity and innate immune responses induced by Nep1-like proteins.** *Plant Cell* 2006, **18**:3721-3744.
31. Ren D, Liu Y, Yang KY, Han L, Mao G, Glazebrook J, Zhang S: **A fungal-responsive MAPK cascade regulates phytoalexin biosynthesis in Arabidopsis.** *Proc Natl Acad Sci U S A* 2008, **105**:5638-5643.
32. Ferrari S, Galletti R, Denoux C, De Lorenzo G, Ausubel FM, Dewdney J: **Resistance to *Botrytis cinerea* induced in Arabidopsis by elicitors is independent of salicylic acid, ethylene, or jasmonate signaling but requires PHYTOALEXIN DEFICIENT3.** *Plant Physiol* 2007, **144**:367-379.
33. Gudesblat GE, Iusem ND, Morris PC: **Guard cell-specific inhibition of Arabidopsis MPK3 expression causes abnormal stomatal responses to abscisic acid and hydrogen peroxide.** *New Phytol* 2007, **173**:713-721.
34. Hamel LP, Nicole MC, Sritubtim S, Morency MJ, Ellis M, Ehltng J, Beaudoin N, Barbazuk B, Klessig D, Lee J *et al.*: **Ancient signals: comparative genomics of plant MAPK and MAPKK gene families.** *Trends Plant Sci* 2006, **11**:192-198.
35. Pitzschke A, Djamei A, Bitton F, Hirt H: **A major role of the •• MEKK1-MKK1/2-MPK4 pathway in ROS signalling.** *Mol Plant* 2009, **2**:120-137.
- Transcriptome analysis of mutants in the MEKK1-MKK1/2-MPK4 pathway revealed a strong overlap and indicated a common deregulation of genes involved in defence mechanisms (salicylic acid related) and ROS homeostasis. In addition, the authors propose MEKK1-controlled signalling pathways that are not mediated through MKK1/MKK2 or MPK4.
36. Teige M, Scheikl E, Eulgem T, Doczi R, Ichimura K, Shinozaki K, Dangl JL, Hirt H: **The MKK2 pathway mediates cold and salt stress signaling in Arabidopsis.** *Mol Cell* 2004, **15**:141-152.
37. Ichimura K, Mizoguchi T, Irie K, Morris P, Giraudat J, Matsumoto K, Shinozaki K: **Isolation of ATMEKK1 (a MAP Kinase Kinase Kinase)-interacting proteins and analysis of a MAP kinase cascade in Arabidopsis.** *Biochem Biophys Res Comm* 1998, **253**:532-543.
38. Brodersen P, Petersen M, Bjorn Nielsen H, Zhu S, Newman MA, Shokat KM, Rietz S, Parker J, Mundy J: **Arabidopsis MAP kinase 4 regulates salicylic acid- and jasmonic acid/ethylene-dependent responses via EDS1 and PAD4.** *Plant J* 2006, **47**:532-546.
39. Miao Y, Laun TM, Smykowski A, Zentgraf U: **Arabidopsis MEKK1 • can take a short cut: it can directly interact with senescence-related WRKY53 transcription factor on the protein level and can bind to its promoter.** *Plant Mol Biol* 2007, **65**:63-76.
- MEKK1 phosphorylates the transcription factor WRKY53 in vitro and phosphorylation could increase DNA-binding activity of WRKY53 in vitro and transcription of a WRKY53 promoter driven reporter gene.
40. Andreasson E, Jenkins T, Brodersen P, Thorgrimsen S, Petersen NH, Zhu S, Qiu JL, Micheelsen P, Rocher A, Petersen M *et al.*: **The MAP kinase substrate MKS1 is a regulator of plant defense responses.** *EMBO J* 2005, **24**:2579-2589.
41. Qiu JL, Fiil BK, Petersen K, Nielsen HB, Botanga CJ, Thorgrimsen S, Palma K, Suarez-Rodriguez MC, Sandbech-Clausen S, Lichota J *et al.*: **Arabidopsis MAP kinase 4 regulates gene expression through transcription factor release in the nucleus.** *EMBO J* 2008, **27**:2214-2221.
- The work reveals the existence of a ternary MKS1-WRKY33-MPK4 complex that depends on the phosphorylation state of MPK4. Active MPK4 phosphorylates MKS1 and leads to the release of WRKY33. Free WRKY33 is then supposed to be responsible for transcriptional activation of downstream genes.
42. Shan L, He P, Li J, Heese A, Peck SC, Nürnberger T, Martin GB, Sheen J: **Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity.** *Cell Host Microbe* 2008, **4**:17-27.
- Two bacterial effectors interact with the BAK1 coreceptor. This interaction interrupts the PAMP-triggered defence responses. AvrPto and AvrPtoB binding to BAK1 seem to be distinct from the AvrPto-Pto interaction, which is responsible for effector-induced immunity.
43. Göhre V, Spallek T, Häweker H, Mersmann S, Mentzel T, Boller T, de Torres M, Mansfield JW, Robatzek S: **Plant pattern-recognition receptor FLS2 is directed for degradation by the bacterial ubiquitin ligase AvrPtoB.** *Curr Biol* 2008, **18**:1824-1832.
44. Zhang J, Shao F, Li Y, Cui H, Chen L, Li H, Zou Y, Long C, Lan L, Chai J *et al.*: **A *Pseudomonas syringae* effector inactivates MAPKs to suppress PAMP-induced immunity in plants.** *Cell Host Microbe* 2007, **1**:175-185.
- HopA1 inactivates MAPKs by removing the phosphate group from phosphothreonine through a unique phosphothreonine lyase activity, thus suppressing the PAMP defence response. The MAPKs MPK3 and MPK6 physically interact with HopA1 indicating that they are direct targets of HopA1.
45. Zhu Y, Li H, Long C, Hu L, Xu H, Liu L, Chen S, Wang DC, Shao F: **Structural insights into the enzymatic mechanism of the pathogenic MAPK phosphothreonine lyase.** *Mol Cell* 2007, **28**:899-913.
46. Li H, Xu H, Zhou Y, Zhang J, Long C, Li S, Chen S, Zhou JM, Shao F: **The phosphothreonine lyase activity of a bacterial type III effector family.** *Science* 2007, **315**:1000-1003.
47. Mazurkiewicz P, Thomas J, Thompson JA, Liu M, Arbibe L, Sansonetti P, Holden DW: **SpvC is a *Salmonella* effector with phosphothreonine lyase activity on host mitogen-activated protein kinases.** *Mol Microbiol* 2008, **67**:1371-1383.
48. Espinosa A, Guo M, Tam VC, Fu ZQ, Alfano JR: **The *Pseudomonas syringae* type III-secreted protein HopPtoD2 possesses protein tyrosine phosphatase activity and suppresses programmed cell death in plants.** *Mol Microbiol* 2003, **49**:377-387.
49. Pfund C, Tans-Kersten J, Dunning FM, Alonso JM, Ecker JR, Allen C, Bent AF: **Flagellin is not a major defense elicitor in *Ralstonia solanacearum* cells or extracts applied to Arabidopsis thaliana.** *Mol Plant Microbe Interact* 2004, **17**:696-706.
50. Citovsky V, Kapelnikov A, Oliel S, Zakai N, Rojas MR, Gilbertson RL, Tzfira T, Loyer A: **Protein interactions involved in nuclear import of the Agrobacterium VirE2 protein in vivo and in vitro.** *J Biol Chem* 2004, **279**:29528-29533.
51. Djamei A, Pitzschke A, Nakagami H, Rajh I, Hirt H: **Trojan horse •• strategy in Agrobacterium transformation: abusing MAPK defense signaling.** *Science* 2007, **318**:453-456 doi: 10.1126/science.1148110.
- The VirE2-interacting protein 1 (VIP1) is shown to be a bZIP transcription factor, which relocates into the nucleus upon phosphorylation via MPK3, where it regulates the expression of PR genes. Bacteria use the MPK3-induced translocation of VIP1 to shuttle the T-DNA-VirE2-VIP1 complex into the nucleus.
52. Tzfira T, Vaidya M, Citovsky V: **Involvement of targeted proteolysis in plant genetic transformation by Agrobacterium.** *Nature* 2004, **431**:87-92.
53. Dardick C, Chen J, Richter T, Ouyang S, Ronald P: **The rice kinase database. A phylogenomic database for the rice kinome.** *Plant Physiol* 2007, **143**:579-586.
54. Popescu SC, Popescu GV, Bachan S, Zhang Z, Gerstein M, Snyder M, Dinesh-Kumar SP: **MAPK target networks in Arabidopsis thaliana revealed using functional protein microarrays.** *Genes Dev* 2009, **23**:80-92.
- A large scale array-based approach was taken to identify potential MAPK substrates.
55. Ritsema T, Joore J, van Workum W, Pieterse CM: **Kinome profiling of Arabidopsis using arrays of kinase consensus substrates.** *Plant Methods* 2007, **3**:3.