



Draft Genome Sequence of *Enterobacter* sp. Sa187, an Endophytic Bacterium Isolated from the Desert Plant *Indigofera argentea*

Feras F. Lafi,^{a,b} Intikhab Alam,^b Rene Geurts,^c Ton Bisseling,^c Vladimir B. Bajic,^b Heribert Hirt,^a  Maged M. Saad^a

King Abdullah University of Science and Technology (KAUST), Biological and Environmental Sciences and Engineering Division (BESE), Thuwal, Kingdom of Saudi Arabia^a; Computational Bioscience Research Center (CBRC), King Abdullah University of Science and Technology (KAUST), Thuwal, Kingdom of Saudi Arabia^b; Department of Plant Sciences, Laboratory of Molecular Biology, Wageningen University, Wageningen, Netherlands^c

ABSTRACT *Enterobacter* sp. Sa187 is a plant endophytic bacterium, isolated from root nodules of the desert plant *Indigofera argentea*, collected from the Jizan region of Saudi Arabia. Here, we report the genome sequence of Sa187, highlighting several genes involved in plant growth-promoting activity and environmental adaptation.

In an effort to explore the microbial diversity of the desert pioneer plants, the Darwin21 project (<http://www.darwin21.net>) has been established. Under the project, extensive microbial isolation from the roots of different desert plants has been conducted. Preliminary results revealed a large diversity of bacterial species with a potential to promote the growth of *Arabidopsis thaliana* plants under different biotic and abiotic stresses. A selected number of these strains were sequenced and characterized as described previously (1, 2). *Enterobacter* sp. Sa187 is an endophytic bacterium isolated from surface-sterilized root nodules formed on roots of the pioneer plant *Indigofera argentea* Burm. f. (*Fabaceae*). Plants were collected from different regions in the Jizan area (16°56.475'N, 42°36.694'E) of Saudi Arabia. Sa187 has been shown to promote plant growth-promoting activities, such as the production of siderophores and indole acetic acid (IAA). Based on the 16S rRNA gene sequence, strain Sa187 is closely related to *E. kobei* CCUG 49023^T and *E. aerogenes* strain KCTC 2190 with 99% sequence similarity (3).

The genomic DNA of Sa187 was extracted using the Qiagen DNeasy blood and tissue kit, following the manufacturer's protocol. The DNA was then sequenced using paired-end Illumina MiSeq, and the library preparation was constructed as described previously (1). Contig assembly was done with SPAdes assembler version 3.6 (4) with a 1-kb contig cutoff size. *De novo* assembly of MiSeq reads for *Enterobacter* sp. Sa187 resulted in 14 contigs with a total length of 4,404,403 bp and a mean contig size of 314,600 bp. The N_{50} was 2,296,004 bp, and the L50 was reached in 1 contig. The G+C content of this draft genome was 56%. MegaBLAST (5) comparison of the Sa187 concatenated contigs against the NCBI reference genome database (<http://www.ncbi.nlm.nih.gov/genome>) revealed the closest relative genomes being *E. sacchari* SP1 with a coverage of 63% and sequence identity of 95% (accession number NZ_CP007215.2) (6). The annotation of *Enterobacter* sp. Sa187 was carried out using the default INDIGO pipeline (7), with the exception of open reading frames (ORFs) predicted by FragGeneScan (8). The annotation of Sa187 resulted in 3,087 ORFs, 9 rRNAs, 75 tRNAs, and 145 ncRNAs.

Received 6 December 2016 Accepted 11 December 2016 Published 16 February 2017

Citation Lafi FF, Alam I, Geurts R, Bisseling T, Bajic VB, Hirt H, Saad MM. 2017. Draft genome sequence of *Enterobacter* sp. Sa187, an endophytic bacterium isolated from the desert plant *Indigofera argentea*. *Genome Announc* 5:e01638-16. <https://doi.org/10.1128/genomeA.01638-16>.

Copyright © 2017 Lafi et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Heribert Hirt, heribert.hirt@kaust.edu.sa.

The annotation predicted a number of siderophore pathway genes such as *entE*, *entC*, *entA*, *entB*, *entF*, as well as *entS*, an MFS transporter of enterobactin. An ABC transporter involved in iron uptake (*sitABCD*) was also found, as well as five copies of the iron complex outer membrane receptor (*fhuA*), and a TonB-dependent outer membrane iron-enterobactin/colicin (*fepA*). Generally, plant growth-promoting rhizobacteria enhance plant growth through the synthesis of IAA from tryptophan via indole pyruvate as the main pathway (9). The Sa187 genome harbors a number of genes involved in this pathway but lacks the gene encoding for indolepyruvate decarboxylase (*ipdC*). Moreover, the Sa187 genome codes for the enzyme tryptophanase (TnaA) (EC: 4.1.99.1), which can transform tryptophan into indole. Further analysis of the genome sequence of Sa187 will provide valuable genetic information to better understand how the strain interacts with different plants.

Accession number(s). The genome of *Enterobacter* sp. Sa187 was deposited at DDBJ/EMBL/GenBank under the accession number [MORB00000000](https://doi.org/10.1128/genomeA.00678-16). The version described in this paper is the first version, MORB00100000.

ACKNOWLEDGMENTS

Genome sequencing was performed at the Bioscience Core Laboratory of the King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia. We are grateful for the use of the Dragon and Snapdragon compute clusters at the Computational Bioscience Research Center (CBRC) of KAUST.

This work was supported by a base fund research grant to H.H. from the King Abdullah University of Science and Technology (KAUST). Computational aspects of this work have been supported by the KAUST Office of Sponsored Research (OSR) under awards URF/1/1976-02 and FCS/1/2448-01 to V.B.B.

REFERENCES

- Lafi FF, Bokhari A, Alam I, Bajic VB, Hirt H, Saad MM. 2016. Draft genome sequence of the plant growth-promoting *Cupriavidus gilardii* strain JZ4 isolated from the desert plant *Tribulus terrestris*. *Genome Announc* 4(4): e00678-16. <https://doi.org/10.1128/genomeA.00678-16>.
- Lafi FF, Ramirez-Prado JS, Alam I, Bajic VB, Hirt H, Saad MM. 2016. Draft genome sequence of *Halomonas elongata* strain K4, an endophytic growth-promoting bacterium enhancing salinity tolerance *in planta*. *Genome Announc* 4(6):e01214-16. <https://doi.org/10.1128/genomeA.01214-16>.
- Shin SH, Kim S, Kim JY, Lee S, Um Y, Oh MK, Kim YR, Lee J, Yang KS. 2012. Complete genome sequence of *Enterobacter aerogenes* KCTC 2190. *J Bacteriol* 194:2373–2374. <https://doi.org/10.1128/JB.00028-12>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
- Zhu B, Chen M, Lin L, Yang L, Li Y, An Q. 2012. Genome sequence of *Enterobacter* sp. strain SP1, an endophytic nitrogen-fixing bacterium isolated from sugarcane. *J Bacteriol* 194:6963–6964. <https://doi.org/10.1128/JB.01933-12>.
- Alam I, Antunes A, Kamau AA, Ba Alawi WB, Kalkatawi M, Stingl U, Bajic VB. 2013. INDIGO—integrated data warehouse of microbial genomes with examples from the Red Sea extremophiles. *PLoS One* 8:e82210. <https://doi.org/10.1371/journal.pone.0082210>.
- Rho MN, Tang HX, Ye YZ. 2010. FragGeneScan: predicting genes in short and error-prone reads. *Nucleic Acids Res* 38:e191. <https://doi.org/10.1093/nar/gkq747>.
- Taghavi S, van der Lelie D, Hoffman A, Zhang YB, Walla MD, Vangronsveld J, Newman L, Monchy S. 2010. Genome sequence of the plant growth promoting endophytic bacterium *Enterobacter* sp. 638. *PLoS Genet* 6:e1000943. <https://doi.org/10.1371/journal.pgen.1000943>.