

AGC kinases in plant development and defense

Heribert Hirt,¹ Ana V. Garcia¹ and Ralf Oelmüller^{2,*}

¹URGV Plant Genomics; INRA-CNRS-University of Evry; Evry, France; ²Friedrich-Schiller-Universität Jena; Institut für Allgemeine Botanik und Pflanzenphysiologie; Jena, Germany

More than 100,000 publications demonstrate that AGC kinases are important regulators of growth, metabolism, proliferation, cell division, survival and apoptosis in mammalian systems.¹ Mutation and/or dysregulation of these kinases contribute to the pathogenesis of many human diseases, including cancer and diabetes. Although AGC kinases are also present in plants, little is known about their functions. We demonstrated that the AGC kinase OXIDATIVE SIGNAL-INDUCIBLE1 (OXI1/AGC2-1) regulate important developmental processes and defense responses in plants. The summary of recent progress also demonstrates that we are only beginning to understand the role of this kinase pathway in plants.

PDK1 Activates AGC Kinases

In mammalian systems, most of the cellular responses to phosphatidylinositol-3-kinase activation and phosphatidylinositol-3,4,5-trisphosphate production are mediated by the activation of a subgroup of Ser/Thr protein kinases termed AGC (cAMP-dependent protein kinase A/cGMP-dependent protein kinase G/protein kinase C), which play essential roles in cell growth, proliferation, survival, metabolism and apoptosis.¹ Many AGC kinases possess a common upstream activator, namely PDK1 (3-phosphoinositide dependent kinase 1), a master kinase which phosphorylates and thus activates the AGC kinases in response to rises in the levels of the second messenger phosphatidylinositol-3,4,5-trisphosphate.^{2,3} Consequently, *pdk1* knock-out mice are embryo-lethal.⁴ PDK1 and AGC kinases

are also present in plants.^{5,6} PDK1 phosphorylates and thus activates AGC kinases such as OXI1 in Arabidopsis⁷ and rice,⁶ or Adi3 (AvrPto-dependent Pto-interacting protein 3) in tomato.⁸ *pdk1* knock-out lines in Arabidopsis and rice are not lethal,⁹ and OXI1 can still be activated in Arabidopsis PDK1-RNAi knock-down lines by stimuli such as H₂O₂ and the pathogen-associated molecular pattern (PAMP) flagellin.¹⁰ Therefore, there are additional stimuli that signal to OXI1 via a PDK1-independent pathway.

PDK1 and its Activation by Phosphatidic Acid

An important second messenger in plant signaling is phosphatidic acid (PA) which can be synthesized either by phospholipase D¹¹ or by a phospholipase C pathway which generates diacylglycerol that is phosphorylated to PA via diacylglycerol kinase.¹² Both lipases are activated in response to many biotic and abiotic stress signals.^{11,12} Recently, it was demonstrated that also the beneficial fungus *Piriformospora indica* is able to stimulate PA synthesis in Arabidopsis.⁹ Therefore, the second messenger PA may integrate various external signals in plants to activate and coordinate appropriate downstream responses. While mammalian PDK1 integrates signals from receptors that stimulate the production of phosphatidylinositol-3,4,5-trisphosphate, the plant PDK1 binds to different signaling lipids, including the second messenger PA.¹³ Thus, although the lipid stimuli are different, animal and plant PDK1 convert phospholipid information into activation of AGC kinases.

Key words: AGC kinases, reactive oxygen species, plant stress, plant microbe interaction, plant pathogen

Submitted: 03/22/11

Accepted: 03/22/11

DOI:

*Correspondence to: Ralf Oelmüller;
Email: b7oera@hotmail.de




Figure 1. The beneficial fungus *Piriformospora indica* stimulates PA synthesis which activates PDK1 and subsequently OXI1 and AGC2-2. On the other hand, OXI1 can also be activated by environmental challenges, including pathogens in a PDK1-independent manner via H₂O₂. PTI1-4 and the MAPKs MPK3 and 6 are downstream factors of OXI1 and regulate the balance between growth and defense responses.

AGC kinases from Arabidopsis

The Arabidopsis genome encodes 39 members of the AGC protein kinase family⁵ and they are involved in various signaling pathways including blue light¹⁴ and auxin signaling.¹⁵⁻¹⁷ Among the AGC kinases, OXI1 was shown to be required for reactive oxygen species (ROS)-mediated responses in Arabidopsis such as root hair elongation and for disease resistance to biotrophic pathogens such as the oomycete *Hyaloperonospora arabidopsidis* and *Pseudomonas syringae* bacteria.^{18,19} The kinase activity of OXI1 itself was induced by H₂O₂, wounding, cellulase and various elicitor treatments mimicking pathogen attack.^{10,18} Furthermore, as *oxi1* mutant plants are impaired in the activation of mitogen-activated protein kinase

(MPK)3 and MPK6 in response to cellular injury and oxidative stress,¹⁸ OXI1 is an upstream regulator of stress-responsive MPKs although its mechanism is still unclear.

In Arabidopsis and rice, kinases of the PTI1 family were identified as interacting partners and kinase targets of OXI1 making them downstream components of OXI1 signaling.^{10,20,21} PTI1 proteins are Ser/Thr protein kinases that share strong sequence identity to tomato PTI1 (Pto-interacting 1). In tomato, PTI1 is phosphorylated by the Ser/Thr kinase Pto conferring resistance to *P. syringae* expressing the effector AvrPto and positively regulates the cell death response triggered by Pto.^{22,23} On the contrary, rice Pti1a inhibits disease resistance and cell death and is negatively regulated by OsPDK1-OsOXI1

signaling cascade in response to ROS and PAMP treatments.^{6,21,24} Interestingly, Arabidopsis Pti1-4 was recently shown to form protein complexes with MPK3 and MPK6 and could therefore mediate OXI1 regulation of the MPKs²⁰ and provides a hypothesis how PTI1 proteins could regulate disease resistance.

The endophytic fungus *P. indica*, a cultivable basidiomycete of Sebaciales, colonizes the roots of many plant species including Arabidopsis.^{25,26} The fungus stimulates growth, biomass and seed production of the hosts²⁵⁻³³ and promotes nitrate and phosphate uptake and metabolism.^{29,34,35} *P. indica* also confers resistance against abiotic^{30,36,37} and biotic stress.^{26,38} In a genetic screen for Arabidopsis mutants which do not respond to *P. indica*, we have identified *OXI1* as the responsible

gene for the growth phenotype induced by *P. indica*.⁹ Interestingly, whereas OXI1 is required for the long-term interaction between Arabidopsis and *P. indica*, the closest homolog of OXI1, AGC2-2, was found to be important for the early steps in the establishment of the symbiosis. This clearly demonstrates that these two AGC2 kinases participate in *P. indica*-induced growth promotion. Since OXI1 can be activated by H₂O₂ and PDK1, we also tested whether mutants in PDK1.1 or PDK1.2, the two PDK1 genes present in the Arabidopsis genome, are defective in the *P. indica*-induced growth phenotype. We found that single mutants still respond to *P. indica* but not *pdk1.1 pdk1.2* double knock-out mutants.⁹ Root colonization by the fungus stimulates PA synthesis in Arabidopsis plants. Phospholipase D (PLD) isoforms synthesize most of the PA in roots in response to stress stimuli. When PA synthesis was reduced by inactivation of PLD α 1 or PLD δ , the *P. indica*-induced growth promotion was compromised.⁹ These results suggest that *P. indica* stimulates growth by PA-mediated activation of PDK1 which subsequently activates OXI1. ROS production is not stimulated and even inhibited by the beneficial fungus and thus does not play a role in activating OXI1.⁹

Open Questions

Based on these observations, plant AGC kinases seem to regulate the interaction with diverse microbes and to be involved in the control of developmental programs, such as cell death, growth stimulation in response to stimuli from endophytic fungi and auxin functions. The model in Figure 1 highlights the present knowledge about OXI1 signaling but also points to the open questions: First, what determines the specificity of OXI1 signaling to a given stimulus? Pathogen attack by biotrophic pathogens such as *H. arabidopsidis* or interaction with the friendly *P. indica* requires the activation of different cellular programs in the plant, and a comparative analysis of the two systems might help to understand how OXI1 discriminates and regulates these processes with different outcomes. In this respect, the discrimination may derive from the

activating stimuli/kinase that may lead to the association of OXI1 with different signaling partners. A recent phosphoproteome analysis of Arabidopsis roots identified differentially phosphorylated proteins in *oxi1* seedlings in response to cellulase treatment.³⁹ Similar approaches could be used to compare potential components of the OXI1 signaling pathway in response to different stimuli. Given that OXI1 shows different expression patterns in roots and aerial tissues,¹⁸ the specificity of OXI1 signaling could also change depending on the plant tissue. Furthermore, in Arabidopsis there are AGC kinases with high sequence homology to OXI1, that also interact with Pti1-4 in yeast²⁰ and the fact that attempts to obtain double homozygous *oxi1 agc2-2* mutants failed⁹ points to certain functional redundancy. Secondly, most biochemical studies on the OXI cascade including phosphorylation and protein/protein interaction assays were performed for defense responses so far but genetic studies and the fact that OXI1 activity is stimulated by growth hormones⁷ suggest that the involvement of this pathway in growth regulation deserves further characterization. The functions of AGC2-2 and PTII-4 have only been analyzed for one model system. Therefore, the relation of OXI1 to other AGC and PTII kinases needs to be investigated in the context of different signaling functions. Considering the intensive work on AGC kinases in the mammalian field,¹ which also lead to medicinal applications, the role of this novel kinase pathway is poorly understood in plants. The phenotypes of Arabidopsis mutants in *agc* kinase genes demonstrate the central role of these protein kinases in development and response to environmental signals.

References

1. Pearce LR, Komander D, Alessi DR. The nuts and bolts of AGC protein kinases. *Nat Rev Mol Cell Biol* 2010; 11:9-22.
2. Bayasas JR. PDK1: The major transducer of PI 3-kinase actions. *Curr Top Microbiol Immunol* 2010; 346:9-29.
3. Mora A, Komander D, van Aalten DM, Alessi DR. PDK1, the master regulator of AGC kinase signal transduction. *Semin Cell Dev Biol* 2004; 15:161-70.
4. Lawlor MA, Mora A, Ashby PR, Williams MR, Murray-Tait V, Malone L, et al. Essential role of PDK1 in regulating cell size and development in mice. *EMBO J* 2002; 21:3728-38.

5. Bögge L, Okrész L, Henriques R, Anthony RG. Growth signaling pathways in Arabidopsis and the AGC protein kinases. *Trends Plant Sci* 2003; 8:424-31.
6. Matsui H, Miyao A, Takahashi A, Hirochika H. *pdk1* kinase regulates basal disease resistance through the OsOxi1-OsPti1a phosphorylation cascade in rice. *Plant Cell Physiol* 2010; 51:2082-91.
7. Anthony RG, Henriques R, Helfer A, Meszaros T, Rios G, Testerink C, et al. A protein kinase target of a PDK1 signaling pathway is involved in root hair growth in Arabidopsis. *EMBO J* 2004; 23:572-81.
8. Devarenne TP, Ekengren SK, Pedley KF, Martin GB. Adi3 is a Pdk1-interacting AGC kinase that negatively regulates plant cell death. *EMBO J* 2006; 25:255-65.
9. Camehl I, Drzewiecki C, Vadassery Y, Shahollari B, Sherameti I, Forzani C, et al. The OXI1 kinase pathway mediates *Piriformospora indica*-induced growth promotion in Arabidopsis. *PLoS Pathogen* 2011; 7:e1002051.
10. Anthony RG, Khan S, Costa J, Pais MS, Bogre L. The Arabidopsis protein kinase PTII-2 is activated by convergent phosphatidic acid and oxidative stress signaling pathways downstream of PDK1 and OXI1. *J Biol Chem* 2006; 281:37536-46.
11. Li M, Hong Y, Wang X. Phospholipase D- and phosphatidic acid-mediated signaling in plants. *Biochim Biophys Acta* 2009; 1791:927-35; Epub 2009 Mar 13; PubMed PMID: 19289179.
12. Arisz SA, Testerink C, Munnik T. Plant PA signaling via diacylglycerol kinase. *Biochim Biophys Acta* 2009; 1791:869-75.
13. Deak M, Casamayor A, Currie RA, Downes CP, Alessi DR. Characterisation of a plant 3-phosphoinositide-dependent protein kinase-1 homologue which contains a pleckstrin homology domain. *FEBS Lett* 1999; 451:220-6.
14. Briggs WR, Christie JM. Phototropins 1 and 2: Versatile plant blue-light receptors. *Trends Plant Sci* 2002; 7:204-10.
15. Christensen SK, Dagenais N, Chory J, Weigel D. Regulation of auxin response by the protein kinase PINOID. *Cell* 2000; 100:469-78.
16. Benjamins R, Quint A, Weijers D, Hooykaas P, Offringa R. The PINOID protein kinase regulates organ development in Arabidopsis by enhancing polar auxin transport. *Development* 2001; 128:4057-67.
17. Robert HS, Offringa R. Regulation of auxin transport polarity by AGC kinases. *Curr Opin Plant Biol* 2008; 11:495-502.
18. Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, Okamoto H, et al. OXI1 kinase is necessary for oxidative burst-mediated signaling in Arabidopsis. *Nature* 2004; 427:858-61.
19. Petersen LN, Ingle RA, Knight MR, Denby KJ. OXI1 protein kinase is required for plant immunity against *Pseudomonas syringae* in Arabidopsis. *J Exp Bot* 2009; 60:3727-35.
20. Forzani C, Carreri A, de la Fuente van Bentem S, Lecourieux D, Lecourieux F, Hirt H. The Arabidopsis protein kinase Pto-interacting 1-4 is a common target of the oxidative signal-inducible 1 and mitogen-activated protein kinases. *FEBS J* 2011; 278:1126-36.
21. Matsui H, Yamazaki M, Kishi-Kaboshi M, Takahashi A, Hirochika H. AGC kinase OsOxi1 positively regulates basal resistance through suppression of OsPti1a-mediated negative regulation. *Plant Cell Physiol* 2010; 51:1731-44.
22. Martin GB, Brommonschenkel SH, Chunwongse J, Frary A, Ganal MW, Spivey R, et al. Map-based cloning of a protein kinase gene conferring disease resistance in tomato. *Science* 1993; 262:1432-6.
23. Zhou J, Loh YT, Bressan RA, Martin GB. The tomato gene Pti1 encodes a serine/threonine kinase that is phosphorylated by Pto and is involved in the hypersensitive response. *Cell* 1995; 83:925-35.

24. Takahashi A, Agrawal GK, Yamazaki M, Onosato K, Miyao A, Kawasaki T, et al. Rice Pti1a negatively regulates RAR1-dependent defense responses. *Plant Cell* 2007; 19:2940-51.
25. Peskan-Berghöfer T, Shahollari B, Giong PH, Hehl S, Markert C, et al. Association of *Piriformospora indica* with *Arabidopsis thaliana* roots represents a novel system to study beneficial plant-microbe interactions and involves early plant protein modifications in the endoplasmic reticulum and at the plasma membrane. *Physiologia Plantarum* 2004; 122:465-77.
26. Oelmüller R, Sherameti I, Tripathi S, Varma A. *Piriformospora indica*, a cultivable root endophyte with multiple biotechnological applications. *Symbiosis* 2009; 49:1-17.
27. Verma S, Varma A. *Piriformospora indica*, gen. et sp. nov., a new root-colonizing fungus. *Mycologia* 1998; 90:896-903.
28. Shahollari B, Vadassery J, Varma A, Oelmüller R. A leucine-rich repeat protein is required for growth promotion and enhanced seed production mediated by the endophytic fungus *Piriformospora indica* in *Arabidopsis thaliana*. *Plant J* 2007; 50:1-13.
29. Sherameti I, Shahollari B, Venus Y, Altschmied L, Varma A, Oelmüller R. The endophytic fungus *Piriformospora indica* stimulates the expression of nitrate reductase and the starch-degrading enzyme glucan-water dikinase in tobacco and *Arabidopsis* roots through a homeodomain transcription factor that binds to a conserved motif in their promoters. *J Biol Chem* 2005; 280:26241-7.
30. Sherameti I, Tripathi S, Varma A, Oelmüller R. The root-colonizing endophyte *Piriformospora indica* confers drought tolerance in *Arabidopsis* by stimulating the expression of drought stress-related genes in leaves. *Mol Plant Microbe Interact* 2008; 21:799-807.
31. Sherameti I, Venus Y, Drzewiecki C, Tripathi S, Dan VM, Nitz I, et al. PYK10, a β -glucosidase located in the endoplasmic reticulum, is crucial for the beneficial interaction between *Arabidopsis thaliana* and the endophytic fungus *Piriformospora indica*. *Plant J* 2008; 54:428-39.
32. Vadassery J, Ranf S, Drzewiecki C, Mithöfer A, Mazars C, Scheel D, et al. A cell wall extract from the endophytic fungus *Piriformospora indica* promotes growth of *Arabidopsis* seedlings and induces intracellular calcium elevation in roots. *Plant J* 2009; 59:193-206.
33. Waller F, Achatz B, Baltruschat H, Fodor J, Becker K, Fischer M, et al. The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance and higher yield. *Proc Natl Acad Sci USA* 2005; 102:13386-91.
34. Shahollari B, Peskan-Berghöfer T, Oelmüller R. Receptor kinases with leucine-rich repeats are enriched in Triton X-100 insoluble plasma membrane microdomains from plants. *Physiologia Plantarum* 2004; 122:397-403.
35. Yadav V, Kumar M, Deep DK, Kumar H, Sharma R, Tripathi T, et al. A phosphate transporter from the root endophytic fungus *Piriformospora indica* plays a role in phosphate transport to the host plant. *J Biol Chem* 2010; 285:26532-44.
36. Baltruschat H, Fodor J, Harrach BD, Niemczyk E, Barna B, Gullner G, et al. Salt tolerance of barley induced by the root endophyte *Piriformospora indica* is associated with a strong increase in antioxidants. *New Phytologist* 2008; 180:501-10.
37. Sun C, Johnson JM, Cai D, Sherameti I, Oelmüller R, Lou B. *Piriformospora indica* confers drought tolerance in Chinese cabbage leaves by stimulating antioxidant enzymes, the expression of drought-related genes and the plastid-localized CAS protein. *J Plant Physiol* 2010; 167:1009-17.
38. Stein E, Molitor A, Kogel KH, Waller F. Systemic resistance in *Arabidopsis* conferred by the mycorrhizal fungus *Piriformospora indica* requires jasmonic acid signaling and the cytoplasmic function of NPR1. *Plant Cell Physiol* 2008; 49:1747-51.
39. Howden AJ, Salek M, Miguet L, Pullen M, Thomas B, Knight MR, et al. The phosphoproteome of *Arabidopsis* plants lacking the oxidative signal-inducible1 (OXI1) protein kinase. *New Phytol* 2010; In press; DOI: 10.1111/j.1469-8137.2010.03582.

©2010 Landes Bioscience.
Do not distribute.